subfossil preservation were not favorable in the early Holocene.

Reconstruction of Climate Variables

Radial growth (tree-ring width) of northern forest-limit pines is limited by concurrent mean July temperatures over the region under study. This particular relationship becomes weaker with increasing distance from the forest limit (Lindholm et al., 2000). In previous studies, approximately 40 to 50% of independent mid-summer temperature variations were captured by transfer models using a single (ultra-long) tree-ring width chronology. Reconstruction of mid-summer temperatures is elongated over the last 7.5 millennia (Fig. 2). Among climate-related variables, North Atlantic Oscillation (NAO) was reconstructed by Lindholm et al. (2001) using a network of pines mainly from Finland and Northwest Russia.

Chronology Signal Strength

The confidence of a climate signal is measured as a function of sample replication and inter-correlation. Extraction of signals of low frequency climate variation from a given long chronological sequence requires a greater number of samples than would be needed to obtain the same level of reliability in signals of high frequency variability. The annual sample replication exceeds thirty trees over 36% of the total chronology length. Approximately

Figure 2: Finnish Lapland pine tree-ring-based mid-summer reconstruction extending back to 5520 BC (data from Eronen et al. 2002; Helama et al. 2002) shown as annual (blue) and centennial (red) anomalies from long-term mean.

13% of the total chronology length is covered by ten or less samples.

Future Prospects

In addition to increasing sample replication, the intra-annual ring-density record can strengthen the desired temperature signal. The maximum latewood density variable could be used along with ring-width variables in transfer functions. Interestingly, chronology could certainly span over the debated 8.2 k event after an elongation of one millennium.

REFERENCES

Eronen, M., Zetterberg, P., Briffa, K., Lindholm, M., Meriläinen, J. and Timonen, M., 2002: The supra-long Scots pine tree-ring record for northern Finnish Lapland – Part 1: chronology construction and initial inferences. *The Holocene* **12**: 673-680.

Helama, S., Lindholm, M., Timonen, M. and Eronen, M., 2002: The supra-long Scots pine tree-ring record for northern Finnish Lapland – Part 2: interannual to centennial variability in summer temperatures for 7500 years. *The Holocene* **12**: 681-687.

Lindholm, M., 1996: Reconstruction of the past climate from ring-width chronologies of Scots pine (*Pinus sylvestris* L.) at the northern forest limit in Fennoscandia. *Publications in Sciences* **40**. PhD thesis. University of Joensuu. 169 p.

Lindholm, M., Eggertsson, Ó., Lovelius, N., Raspopov, O., Shumilov, O. and Läänelaid, A., 2001: Growth indices of North European Scots pine record the seasonal North Atlantic Oscillation. *Boreal Environment Research* **6**: 275-284.

Lindholm, M., Lehtonen, H., Kolström, T., Meriläinen, J., Eronen, M. and Timonen, M., 2000: Climatic signals extracted from ring-width chronologies of Scots pine from the Northern, Middle and Southern parts of the boreal forest belt in Finland. *Silva Fennica* **34**: 317-329.

For full references please consult:

www.pages-igbp.org/products/newsletters/ref2003_2.html

Climate Reconstruction From Peatlands

DAN CHARMAN¹ AND MARKKU MÄKILÄ²

¹Dept Geographical Sciences, University of Plymouth, Plymouth, UK; dcharman@plymouth.ac.uk,

²Geological Survey of Finland, Espoo, Finland; markku.makila@gsf.fi

Peatlands occur throughout the mid to high latitudes of the northern hemisphere, with huge expanses in low-relief areas of the oceanic and boreal zones. Smaller areas occur in mountainous regions throughout the world, in oceanic temperate zones in the southern hemisphere, and in the tropics. The vast majority of the peat is Holocene in age, often forming

long, continuous sequences from the early to mid Holocene up to the present day. Accumulation rates are highly variable, but 0.5-1 cm per decade is typical for raised bogs in Europe. For example, the long-term average peat accumulation rate for 210 raised bog profiles in Finland is 0.6 cm per decade. The deposits are mainly autochthonous and relatively straightforward to date

with radiocarbon, especially where mosses dominate the peat. Therefore, peatlands potentially provide a widespread source of well-dated paleoenvironmental data, covering most of the Holocene at decadal resolution. Despite this, their use for paleoclimatic reconstructions has still been limited to a relatively small number of studies in a few main locations. Recent innovations

Figure 1: Vegetation and surface patterns (hummocks and hollows) of a raised bog (Kilpisuo) in southern Finland, one of the sites used in studies of carbon accumulation and climate change (Mäkilä, 2001; Mäkilä, et al., 2001).

in techniques applicable to peatlands, better chronological precision of the record, and improvements in our understanding of how peatlands record climate history, mean that the time is now ripe to begin to exploit these deposits more fully. Here, we outline some of the main approaches, current research and ways in which peat records may be developed in the future to meet the needs of Holocene paleoclimate research.

Approaches to Reconstruct Past Climates Using Peat Deposits:

• *Records of changing surface wetness.* Most techniques aim to reconstruct past surface wetness as a proxy for precipitation-evaporation (PE) balance. These include plant macrofossil analysis, peat humification and testate amoebae analysis. The surface wetness of all peatlands is dependent to some extent on PE but in fact ombrotrophic peatlands (raised bogs or watershed blanket mires) are the only sites where there is no influence of surface runoff or groundwater whatsoever, and precipitation and evapotranspiration are the only components of water balance.

• *Records of temperature change.* Recent work has shown the considerable potential of proxies that record temperature directly. Principal amongst these are species-specific and compound-specific isotopic analysis (e.g. Pendall et al., 2001; Xie

et al., 2000), and mercury analysis (Martinez-Cortizas et al. 1999).

• *Records of changing mass and carbon accumulation rates.* Increased accumulation rates may reflect periods of more positive PE balance (Mäkilä et al., 2001; Mäkilä, 2001) (Fig. 1).

• *The timing and rate of spread of peat initiation.* The most successful of these studies have been in western Canada using large data sets from very wide geographic areas, where peat initiation data have been integrated with other paleoclimatic indices and climate models (Gajewski et al., 2000), and cyclicities similar to those found in ocean and ice core records have been shown (Campbell et al., 2000).

Reconstruction of past surface wetness changes is the basis of many existing records, including some

that use multiple records from several sites in the same climate area to provide composite records (Fig. 2). Many of the records from north-west Europe display cyclicities with periodicities of varying lengths including possible solar-related cycles (Chambers and Blackford, 2001) and links to ocean variability on longer time-scales (Hughes et al. 2000).

One key issue in current research is to resolve how long-term changes in surface wetness relate to seasonal and annual climate variability. Conceptual process models suggest that surface wetness changes reflect some combination of temperature, precipitation, humidity and wind speed. The precise nature of this relationship is unknown and is likely to be different in different climate zones. Variability in surface wetness is probably primarily related to summer climate because peatlands are either saturated or frozen in the winter months. Comparisons with documentary data suggest that both precipitation and temperature may have determined surface wetness in northern England over the past 900 years (Barber et al., 1994) (Fig. 2). Comparisons of high-resolution records with instrumental climate data confirm that summer climate is the main driver of wetness changes but that the relationship with temperature and precipitation probably varies in different locations across Europe (Table 1). Under a prevailing oceanic climate such as that in northern England, summer precipitation may be the main factor reflected in reconstructed water table variability. Under a more continen-

Table 1: Linear correlation coefficients between reconstructed water table records from two European raised bogs (based on testate amoebae analysis) and instrumental climate data. The Butterburn Flow record covers AD 1800-1990 and is based on decadal averages ($n=20$ for temperature, $n=17$ for precipitation). The Männikjärve record covers AD 1951-1995 and is based on 5-year averages ($n=9$). * $p<0.05$. From Charman et al. (submitted).

Climate variable	Butterburn Flow England	Männikjärve bog Estonia
Temperature		
Summer	-0.046	*-0.698
Growing season	-0.127	-0.176
Mean annual	0.026	-0.060
Precipitation		
Summer	*0.458	0.437
Growing season	0.106	0.377
Mean annual	0.188	*0.585

tal regime such as that in northern Europe, winter precipitation stored as snow can also have an effect on reconstructed water tables. Snow melts rapidly in springtime filling hollows and causing heavy floods. Summer temperature also exerts a much stronger desiccating effect in more continental sites, affecting the moisture balance in surface peats directly. New and improved analytical approaches and better understanding of the peat-climate system now suggest a number of ways forward for maximizing the use of the peat record.

Key Areas for the Improvement of Climate Reconstructions:

- *Understanding the peatland-climate relationship.* Calibration of records with instrumental data is critical in deriving empirical relationships between changes in proxy data and climate parameters (see above). Development of physical process models from hydrological, biological and meteorological data would lead to greater confidence in empirical relationships and extrapolating calibrated records over pre-instrumental time periods. Comparisons with other independent climate proxies will also help test hypothesized surface wetness-climate links.

- *Further development of new proxy climate techniques.* Isotopic, compound-specific and mercury analyses are now producing exciting results. These approaches need to be explored more extensively.

- *Establishment of additional records from different climatic regimes.* Much existing work has been based in oceanic temperate regions. Additional work on continental, alpine and higher latitude areas is needed to fully exploit the potential peatland archive. Much of this work is already underway, as is reflected in recent results from central Europe (Mitchell et al., 2001), continental North America (Booth, 2002), South America (Pendall, 2001) and New Zealand (e.g. McGlone and Wilmshurst, 1999; Wilmshurst et al., 2002).

- *Improving chronological control.* Existing comparisons between

Figure 2: Mean annual water table changes for the last 1500 years, reconstructed from testate amoebae analysis of three peat profiles in northern England (c), compared with documentary index of summer wetness and winter severity (a) and ice accumulation record from GISP2 (d). Instrumental records of mean annual temperature and precipitation are also shown (b). All data are 100-year running averages. Raw documentary and water table data are also shown. Fluctuations in the reconstructed water tables, documentary indices and ice accumulation show similar trends over this time period. See Charman and Hendon (2000).

peatland records and other proxies such as speleothem, ice core and ocean records (e.g. Charman and Hendon, 2000; Charman et al., 2001; Pendall et al., 2001) are limited by chronological uncertainty. Calibrated radiocarbon ages still often mean possible 2σ errors of ± 200 years—critical in trying to establish whether climate variability on centennial scales is in or out of phase. Wiggle-matched radiocarbon ages (Mauquoy et al., 2002) and tephrochronology (e.g. Van Den Bogaard and Schmincke, 2002) both have the potential to significantly improve chronologies and the precision of correlations with other records.

Conclusions:

One of the main HOLIVAR themes is the comparison of different proxy climate records—with all the attendant problems of different chronologies, climate parameters recorded and geographical coverage. With the development of a wider range of techniques and a better understanding of the proxy-climate relationship linked to improved chronologies,

peatland paleoclimate records can now be fully integrated with other Holocene paleoclimate reconstructions.

REFERENCES

- Charman, D.J., Brown, A.D., Hendon, D., Kimmel, A. and Karofeld, E., Testing the relationship between Holocene peatland palaeoclimate reconstructions and instrumental data. *Quaternary Science Reviews* (submitted).
- Charman, D.J. and Hendon, D., 2000: Long-term changes in soil water tables over the past 4500 years: Relationships with climate and North Atlantic atmospheric circulation and sea surface temperature. *Climatic Change* **47**: 45-59.
- Charman, D.J., Caseldine, C., Baker, A., Gearey, B.R. and Hatten, J., 2001: Palaeohydrological records from peat profiles and speleothems in Sutherland, northwest Scotland. *Quaternary Research* **55**: 223-234.
- Mäkilä, M., 2001: Climate in relation to carbon accumulation on Kilpisuo, a raised bog in southern Finland. In: Vasiliev, S. V., Titlyanova, A. A. and Velichko, A. A. (eds.) *West Siberian peatlands and carbon cycle: Past and present. Proceedings of an International Symposium Noyabrsk*. pp.38-40.
- Mäkilä, M., Saarnisto, M. and Kankainen, T., 2001: Aapa mires as a carbon sink and source during the Holocene. *Journal of Ecology* **89**: 589-599.

For full references please consult:
www.pages-igbp.org/products/newsletters/ref2003_2.html