Figure 2: **A)** Detrended microscopic charcoal (green line) and oxygen isotope record (purple line) for Eski Acigöl Lake (Turkey). Correlation analysis (gray bars) shows statistically significant correlations between microscopic charcoal influx (particles $\text{cm}^{-2} \text{a}^{-1}$) and $\delta^{18}\text{O}$ (‰) for most of the record (modified from Turner et al., 2008); **B)** Lago dell'Accesa (Tuscany, Italy) lake-level fluctuations (blue line; modified from Magny et al., 2007) and the fire-return interval (red line) reconstructed from fire-event detection based on sedimentary macroscopic charcoal (modified from Vannière et al., 2008). Peaks in the fire-return interval curve correspond with high lake-level stands before 4 cal ka BP.

across the Mediterranean (Carrión et al., 2003, 2007; Sadori et al., 2008).

Conclusion

The paleofire record from the Mediterranean is paradoxical. Climatic variations have certainly acted as one of the main pacemakers of fire regimes, particularly in the first half of the Holocene. Under different climate conditions (e.g., seasonality of precipitation), the southern and northern Mediterranean may have been

differentially impacted by fire. Similarly, human actions (e.g., directly via ignition or indirectly via fuel management) have both increased and decreased fire activity during the Holocene. Increased sedimentary charcoal influx is often associated with pre- and proto-historic forest clearance but in the late Holocene, wildfire frequency often reached a maximum during phases of land abandonment and secondary scrub-woodland development, e.g., during the last century in much of Medi-

terranean Europe. Even apparently well-established relationships, such as evergreen oaks being favored by fire, turn out to be wrong when viewed over decadal to centennial timescales. These complex long-term responses are significant in the context of increasing aridity and warming, as well as major regional land-use changes linked to agricultural and tourism development around the Mediterranean Sea. Understanding them will help us to better manage and preserve one of the most fire-prone regions of the world, characterized by extraordinary plant diversity.

Data

The data are submitted to the Global Palaeofire Working Group Database (<http://www.gpwg.org>) and are available by contacting B. Vannière (boris.vanniere@univ-fcomte.fr).

Acknowledgements

We are greatly indebted to M. Magny, M. Power, W. Tinner and R. Turner who have contributed to these results and for enriching discussions about paleofire. Many thanks also to all contributors to the Mediterranean part of the Global Charcoal Database.

References

- Colombaroli, D., Tinner, W., van Leeuwen, J.F.N., Noti, R., Vescovi, E., Vannière, B., Magny, M., Schmidt, R. and Bugmann, H., 2009: Response of broad-leaved evergreen Mediterranean forest vegetation to fire disturbance during the Holocene: insights from the peri-Adriatic region, *Journal of Biogeography*, **36**: 314–326.
- Colombaroli, D., Vannière, B., Chapron, E., Magny, M. and Tinner, W., 2008: Fire-vegetation interactions during the Mesolithic-Neolithic transition at Lago dell'Accesa, Tuscany, Italy, *The Holocene*, **18**: 679–692.
- Turner, R., Roberts, N. and Jones, M.D., 2008: Climatic pacing of Mediterranean fire histories from lake sedimentary microcharcoal, *Global and Planetary Change*, **63**: 317–324.
- Vannière, B., et al., in press: Circum-Mediterranean fire activity and climate changes during the mid Holocene environmental transition (8500–2500 cal yr BP), *The Holocene*.
- Vannière, B., Colombaroli, D., Chapron, E., Leroux, A., Tinner, W. and Magny, M., 2008: Climate versus human-driven fire regimes in Mediterranean landscapes: the Holocene record of Lago dell'Accesa (Tuscany, Italy), *Quaternary Science Reviews*, **27**: 1181–1196.

For full references please consult:

http://www.pages-igbp.org/products/newsletters/ref2010_2.html

Tropical fire ecology across the African continent: A paleoecological perspective

DANIELE COLOMBAROLI^{1,2} AND DIRK VERSCHUREN²

¹Institute of Plant Sciences and Oeschger Center for Climate Change Research, University of Bern, Switzerland; daniele.colombaroli@ips.unibe.ch

²Limnology Unit, Department of Biology, Ghent University, Belgium

High-resolution charcoal records from African lake sediments provide new insights for longstanding research questions on fire-climate-human interactions in tropical and subtropical ecosystems.

Every year, the tropics experience more fire than any other region in the world (Fig. 1). Tropical grassland (savannah) fires are the dominant source of carbon from biomass burning and provide more than 60% of the global total (Mouillot and Field, 2005). However, much of what is known about tropical fire ecology is based on

monitoring programs started within the last decade, with highly fragmentary historical data extending only to the early 20th century. These data do not allow us to assess whether recent trends in fire frequency and magnitude are unusual in the context of natural long-term ecosystem dynamics (Gillson and Willis, 2004).

Natural or anthropogenic fire regime?

Seventy percent of tropical and subtropical areas worldwide are considered to have ecologically degraded fire regimes (Shlisky et al., 2007). In Africa, the recent increase in fire frequency is attributed to human ecosystem disturbance associated

with intensifying agriculture (Fig. 2; Davidson et al., 2003). For instance, in the low-land rainforests of West Africa and in moist montane forests at higher elevations in East Africa, natural fire is uncommon (Goldammer, 1990). Yet widespread clearance of natural vegetation has converted large, formerly forested areas into highly flammable grasslands (Roberts, 2000; Goldewijk, 2001). Studies of global-scale patterns in historical land use (e.g., Archibald et al., 2005) tend to assume that human impact on tropical African ecosystems was limited before 1700 AD because population densities of indigenous people were low (Goldewijk, 2001; Ramankutty and Foley, 1999). This perspective contrasts with archaeological and paleoecological evidence that indicates that anthropogenic forest clearance in parts of East Africa started at least 2.5 ka ago, in association with the introduction of iron smelting technology (Robertshaw and Taylor, 2000). Other authors suggest that humans have altered African forest ecosystems over a longer time (Willis et al., 2004). If increasing fire activity during the Holocene was indeed related to intensifying human impact (Lejju et al., 2005; Ryner et al., 2008), the timing and extent to which humans altered local ecosystems varied regionally and among ecosystems. For instance, recent studies indicate that deforestation associated with sedentary agriculture started only ~0.35 ka ago in the moist highlands of central Kenya (Lamb et al., 2003) but at least ~0.8–1 ka ago in sub-humid western Uganda (Ssemmanda et al., 2005; Russell et al., 2009). In drier environments, agricultural activity often began ~0.12 ka ago, during colonial times, yet landscapes may have been significantly modified by pastoralist cultures well before then. Detailed charcoal studies with adequate spatial coverage are needed to determine whether current fire regimes are within the range of historic variability (Willis and Birks, 2006), or whether fire frequency has increased in response to the different types and intensities of human impact associated with pastoralist and agriculturalist societies.

Fire regime response & feedback to past climate variability

High-resolution charcoal studies have shown how fire can be a “catalyst” for climate-change effects on vegetation. For instance, moist conditions limit fire to spread in present tropical forests, but during drier periods in the past wildfire was likely more common, causing changes in ecosystem structure and degradation (e.g., Willis and Birks, 2006; Bush et al.,

2008; Fig. 2). Climate-proxy information from the sediments of East African lakes document major variations in moisture balance. For example, in the late 18th century, an episode of severe drought completely desiccated all but one lake in the Eastern Rift Valley of Kenya, south of Lake Turkana (Verschuren, 2004; Bessems et al., 2008). In the last few millennia, century-long periods of both significantly drier and wetter conditions than today have occurred over most of equatorial East Africa. There have also been periods (e.g., from ~1500 to 1750 AD) when climate was unusually dry in the normally sub-humid western parts of the region while remaining unusually wet in semi-arid regions further east (Verschuren et al., 2000; Russell and Johnson, 2007). Research documenting the response of terrestrial ecosystems to this climate variability reveals the high sensitivity of vegetation transition zones, such as the forest/savannah ecotone, to even modest decadal-scale variations in rainfall (Lamb et al., 2003; Ngomanda et al., 2007). Additional charcoal records of high temporal resolution are needed to show how fire regimes have responded to contrasting climate trends at the regional scale (Fig. 1b).

Tropical ecosystems resilience to fire

In the seasonally dry climate regime prevailing throughout most of East Africa, fire is the dominant direct control on vegetation distribution (Bond et al., 2005; Gillson and Duffin, 2007) but natural fire frequency decreases from semi-arid central and eastern Kenya to sub-humid western Uganda. How does fire control the landscape-scale ecotone between savannah and forest? Recent studies postulate that tropical rainforests and grass savannah may exist as “stable states” in which a grass-dominated ecosystem is maintained by frequent fires while a tree-dominated ecosystem helps create a wet microclimate and low ground cover/fuel load that limits fire (Sankaran et al., 2005; Gillson and Duffin, 2007; Gillson, 2008). A shift from small patchy fires set by indigenous peoples to the larger fires characteristic of European land management has strongly altered the savannah-forest ecotone, by favoring highly flammable annual grasses. Thus, by increasing the flammability of grass communities, this historical change in fire management may have caused a positive feedback with fire (Cochrane, 2009). This hypothesis needs to be rigorously tested

Figure 1: **A)** Satellite image of world fire activity detected by MODIS in the early spring of 2010 (red dots, data from Fire Information for Resource Management System FIRMS, <http://maps.geog.umd.edu/firms>). **B)** Worldwide geographical distribution of paleoecological fire-regime records currently in the Global Charcoal Database (modified from Power et al., 2008). Despite the great fire activity in tropical and subtropical ecosystems, few paleofire records are available from regions such as Africa. Dashed lines delimit the tropical region, bounded by latitudes 23.5°N and S.

Figure 2: **A)** Human-set fire in a rural area close to Mozambique border. Here, fire is used to clear land for agriculture and livestock ranching. **B)** Wooded savannah stand in Kruger National Park (South Africa). During the dry season (Apr-Oct), most of the biomass dries, increasing the fire hazard. (Photos by D. Colombaroli). **C)** Pollen and charcoal record from Malahlapanga Lake (Kruger National Park) showing the transition from grassland to a fire-maintained wooded savannah after 1100 AD (modified from Gillson and Ekblom, 2009). The authors suggested that increased regional rainfall promoted biomass production (i.e., wooded savannah), allowing fire activity to increase.

on long timescales and in multiple regions to determine whether this feedback is characteristic of presently highly disturbed conditions, or whether it also occurred during natural cycles of long-term hydrological change. Paleoinformation on the resilience of tropical moist forests to occasional fire and, specifically, the rate at which rainforests recover from destructive fire would also be highly instructive for future conservation (Cochrane, 2003). New reconstructions of past fire regimes based on fossil charcoal analysis that quantify the local frequency of fire (e.g., Whitlock and Larsen, 2001; Gavin et al., 2006; Higuera et al., 2008), combined with modern calibration studies (Duffin et al., 2008), should reveal how African ecosystems respond to fire variability at decadal to century timescales.

Research outlook

Coupled atmosphere/ocean/biosphere climate models project future temperatures across tropical Africa to increase from 0.2 to 0.5°C per decade, and pre-

cipitation in East Africa to increase during the short rainy season (Northern Hemisphere winter) and decrease during the main rainy season (Northern Hemisphere spring) (Hulme et al., 2001; IPCC, 2007). In addition, changes in the teleconnected El Niño/Southern Oscillation (ENSO) are projected to cause pronounced drought in some regions and increased risk of flooding in others (Wara et al., 2005). If mean annual precipitation over East Africa does increase (IPCC, 2007), its beneficial effect on forest ecosystems will likely be lost in areas with frequent anthropogenic fires and increasing demographic pressure. Insights into how Africa's forest and savannah ecosystems will respond to the multiple stressors of future global climate change requires an understanding of past ecosystem responses to large-magnitude environmental changes. Currently still very rare in Africa and elsewhere in the tropics, high-resolution paleoecological records of (pre-)historical human impact, fire and vegetation can provide such holistic information.

Acknowledgements

We thank Lindsey Gillson for useful comments and Mitchell J. Power for the global map of sites in the Global Charcoal Database. The project "Fire, climate change and human impact in tropical ecosystems: paleoecological insights from the East African region" is funded by the Swiss National Science Foundation (FNS "Ambizione" Project no. PZ00P2_126573).

References

- Cochrane, M.A., 2003: Fire science for rainforests, *Nature*, **421**: 913–919.
- Duffin, K.I., Gillson, L. and Willis, K.J., 2008: Testing the sensitivity of charcoal as an indicator of fire events in savanna environments: quantitative predictions of fire proximity, area and intensity, *The Holocene*, **18**: 279–291.
- Gillson, L. and Ekblom, A., 2009: Resilience and thresholds in savannas: nitrogen and fire as drivers and responders of vegetation transition, *Ecosystems*, **12**: 1189–1203.
- Sankaran, M., Hanan, N. and Scholes, R., 2005: Determinants of woody cover in African savannah, *Nature*, **438**: 846–849.
- Willis, K.J., Gillson, L. and Brncic, T.M., 2004: How "virgin" is virgin rainforest? *Science*, **304**: 402–403.

For full references please visit:

http://www.pages-igbp.org/products/newsletters/ref2010_2.html