

(e.g., Kokfelt et al., submitted), to fully understand permafrost dynamics and the relationship with atmospheric methane concentrations over longer timescales. It is clear from paleorecords that peatland permafrost has expanded and contracted over the Holocene at different times in different places (e.g., Vardy et al., 2005; Oksanen, 2006). Although the impacts of Holocene peatland expansion on atmospheric methane are now being explored (Smith et al., 2004; Korhola et al., 2010; Beilman et al., this issue), the implications of Holocene permafrost variability on past global methane concentrations have not yet been assessed. The sparse data on

contemporary methane emissions show that peatlands with and without permafrost differ significantly in their functioning. More continuous measurements are required to document ongoing changes. Modern process models of carbon dynamics linked to paleoreconstructions of permafrost could produce critical insights into long-term role and functioning of northern peatlands in the global carbon cycle.

References

Johansson, T., Malmer, N., Crill, P.M., Friborg, T., Akerman, J.H., Mastepanov, M. and Christensen, T.R., 2006: Decadal vegetation changes in a northern peatland, greenhouse gas fluxes and net radiative forcing, *Global Change Biology*, **12**: 2352-2369.

Mastepanov, M., Sigsgaard, C., Dlugokencky, E.J., Houweling, S., Strom, L., Tamstorf, M.P. and Christensen, T.R., 2008: Large tundra methane burst during onset of freezing, *Nature*, **456**: 628-631.

McGuire A.D., Anderson L.G., Christensen T.R., Dallimore S., Guo L., Hayes D.J., Heimann M., Lorenson T.D., Macdonald R.W. and Roulet N., 2009: Sensitivity of the carbon cycle in the Arctic to climate change, *Ecological Monographs*, **79**: 523-555.

Schuur, E.A.G., Vogel, J.G., Crummer, K.G., Lee, H., Sickman, J.O. and Osterkamp, T.E., 2009: The effect of permafrost thaw on old carbon release and net carbon exchange from tundra, *Nature*, **459**: 556-559.

Walter, K.M., Edwards, M.E., Grosse, G., Zimov, S.A. and Chapin, F.S., 2007: Thermokarst lakes as a source of atmospheric CH₄ during the last deglaciation, *Science*, **318**: 633-636.

For full references please consult:
http://www.pages-igbp.org/products/newsletters/ref2010_1.html





New on the PAGES Bookshelf

Two new volumes of the Springer *Developments in Paleoenvironmental Research* book series are now available. For more information on both titles visit the Springer website www.springer.com/series/5869?detailsPage=titles&changeHeader=true

Past Climate Variability in South America and Surrounding Regions: From the Last Glacial Maximum to the Holocene

Editors: F. Vimeux, F. Sylvestre and M. Khodri
 2009, ISBN: 978-90-481-2671-2

This book covers tropical, temperate and high latitudes climate variability, and discusses climate mechanisms, specifically the low to high latitude teleconnections in South America.



The South China Sea: Paleooceanography and Sedimentology

Editors: P. Wang and Q. Li
 2009, ISBN 978-1-4020-9744-7

This book is the first synthesis of sedimentary geology and paleoceanography of the South China Sea on the basis of extensive industrial explorations and scientific expeditions.

Climate of the last million years: New insights from EPICA and other records

Editors: H. Fischer, V. Masson-Delmotte, C. Waelbroeck and E.W. Wolff
Quaternary Science Reviews, 2010, 29: ISSN 0277-3791

This special issue of *Quaternary Science Reviews* stems from the EPICA 2008 Open Science Conference (Venice, Italy), which brought together the international ice core community and a large group of researchers from the marine and terrestrial realm, and climate and ice sheet modeling disciplines. This issue therefore represents a compilation of papers on current advances in the science of this period.



For more information, visit the journal website at www.sciencedirect.com/science/journal/02773791